

## EDUCATION COMMITTEE UPDATE By Phil Losavio

### Hyoid Suspension Position Statement

October 2024

Hyoid myotomy/suspension (HS), whether performed separately or combined with other upper airway procedures, **is considered effective and non-investigational** as part of the comprehensive surgical management of symptomatic adult patients with mild obstructive sleep apnea (OSA) and adult patients with moderate and severe OSA assessed as having tongue base or hypopharyngeal obstruction.

HS has been studied rigorously in many patients since it was originally described in 1984. Original research studies include a total of nearly one-thousand patients with treatment results reported in the medical literature.<sup>1-21</sup> Studies have included different variations of the technique such as thyrohyoidpexy and hyoid myotomy with mandibular suspension. Most studies were conducted as part of a multilevel surgery approach in conjunction with pharyngeal and/or nasal surgery. Surgical success as defined by the Sher criteria (AHI less than 20 with a 50% reduction) demonstrates a range of results from 17-90%.<sup>23</sup> 16 studies contain available data regarding this surgical success criterion with 9 of 16 showing success rates greater than 60%. The majority of this past research was based on prospective cohort and retrospective cases series. HS is well tolerated with few complications. The most notable being transient dysphagia, hematoma/seromas, and hardware related complications.<sup>7,14,21</sup>

More recent literature with stricter and more uniform methodologies demonstrates even higher success. Panah et al<sup>20</sup> and Van Tassel et al<sup>21</sup> published more recent studies in 2023. Panah and colleagues conducted an industry sponsored randomized control trial where they divided a group of forty-eight patients into two treatment cohorts that received uvulopalatopharyngoplasty (UP3) and tongue base radiofrequency either with or without modified thyrohyoid suspension. Mean BMI was less than 28. Success was much greater in the hyoid group (75%) compared to the group without hyoid surgery (41.7%) as part of the multilevel protocol ( $p = 0.021$ ). Van Tassel and colleagues investigated a series of thirty-nine patients undergoing hyoid myotomy and suspension combined with uvulopalatopharyngoplasty. 76.9% patients had surgical success with a mean reduction of AHI from 49.9 +/- 25.6 to 15.4 +/-14.9 ( $p < .001$ ). For HS performed as a standalone intervention, there is less overall evidence, but the results are consistent with the additive efficacy effect seen when HS is utilized as part of combined multi-level or staged sequential approach to treatment. UP3 should be considered in conjunction either at the same time or in staged sequence depending on clinical circumstances and best judgement of the surgeon.

Hyoid suspension is safe and effective and should be considered a part of a comprehensive approach in the medical and surgical management of symptomatic adult patients with mild OSA and adult patients with moderate and severe OSA who have evidence of tongue base or hypopharyngeal obstruction. A comprehensive medical evaluation taking into account diagnostic workup (e.g., history and physical, drug-induced sleep endoscopy), medical risk and individual patient factors is needed to make a final determination regarding the appropriate indication for surgery.



## REFERENCES

1. R. Riley, C. Guilleminault, N. Powell, S. Derman, Mandibular Osteotomy and Hyoid Bone Advancement for Obstructive Sleep Apnea: A Case Report, *Sleep*, Volume 7, Issue 1, September 1984, Pages 79–82, <https://doi.org/10.1093/sleep/7.1.79>
2. Riley RW, Powell NB, Guilleminault C. Obstructive sleep apnea and the hyoid: a revised surgical procedure. *Otolaryngol Head Neck Surg.* 1994 Dec;111(6):717-21. doi: 10.1177/019459989411100604. PMID: 7991249.
3. Krespi Y, Kacker A. Hyoid suspension for obstructive sleep apnea. *Oper Tech in Otol-Head and Neck Surg.* 2002;13(2):144-149.
4. Neruntarat C. Hyoid myotomy with suspension under local anesthesia for obstructive sleep apnea syndrome. *Eur Arch Otorhinolaryngol.* 2003 May;260(5):286-90.
5. Vilaseca I, Morelló A, Montserrat JM, Santamaría J, Iranzo A. Usefulness of uvulopalatopharyngoplasty with genioglossus and hyoid advancement in the treatment of obstructive sleep apnea. *Arch Otolaryngol Head Neck Surg.* 2002;128(4):435-40.
6. den Herder C, van Tinteren H, de Vries N. Hyoidthyroidpexia: a surgical treatment for sleep apnea syndrome. *Laryngoscope.* 2005 Apr;115(4):740-5.
7. Bowden MT, Kezirian EJ, Utley D, Goode RL. Outcomes of hyoid suspension for the treatment of obstructive sleep apnea. *Otolaryngol Head Neck Surg.* May 2005; 131(5):440-445.
8. Baisch A, Maurer JT, Hormann K. The effect of hyoid suspension in a multilevel surgery concept for obstructive sleep apnea. *Otolaryngol Head Neck Surg.* 2006 May;134(5):856-61.
9. Verse T, Baisch A, Maurer JT, Stuck BA, Hormann K. Multilevel surgery for obstructive sleep apnea: short-term results. *Otolaryngol Head Neck Surg.* 2006 Apr;134(4):571-7.
10. Benazzo M, Pagella F, Matti E, Zorzi S, Campanini A, Frassinetti S, Montevecchi F, Tinelli C, Vicini C. Hyoidthyroidpexia as a treatment in multilevel surgery for obstructive sleep apnea. *Acta Oto-Laryngologica*, 2008; 128: 680-684.
11. Gillespie MB, Ayers CM, Nguyen SA, Abidin MR. Outcomes of hyoid myotomy and suspension using a mandibular screw suspension system. *Otolaryngol Head Neck Surg.* Feb 2011; 144(2):225-229.
12. Karataylı-Özgürsoy S, Demireller A. Hyoid suspension surgery with UPPP for the treatment of hypopharyngeal airway obstruction in obstructive sleep apnea. *Ear Nose Throat J.* 2012 Aug;91(8):358-64.
13. Berg EE, Bunge F, Delgado JM. Multilevel treatment of moderate and severe obstructive sleep apnea with bone-anchored pharyngeal suspension sutures. *Ear Nose Throat J.* 2013 Aug;92(8):E1.
14. Canzi P, Berardi A, Tinelli C, Montevecchi F, Pagella F, Vicini C, Benazzo M. Thirteen Years of Hyoid Suspension Experience in Multilevel OSAHS Surgery: The Short-Term Results of a Bicentric Study. *Int J Otolaryngol.*
15. Piccin O, Scaramuzzino G, Martone C, Marra F, Gobbi R, Sorrenti G. Modified hyoid suspension technique in the treatment of multilevel related obstructive sleep apnea. *Otolaryngol Head Neck Surg.* 2014 Feb;150(2):321-4. doi:
16. Verse T, Wenzel S, Brus J. Multi-level surgery for obstructive sleep apnea. Lingual tonsillectomy vs. hyoid suspension in combination with radiofrequency of the tongue base. *Sleep Breath.* 2015 Dec;19(4):1361-6. doi: 10.1007/s11325-015-1241-8.
17. Ong AA, Buttram J, Nguyen SA, Platter D, Abidin MR, Gillespie MB. Hyoid myotomy and suspension without simultaneous palate or tongue base surgery for obstructive sleep apnea. *World Journal of Otorhinolaryngology- Head and Neck Surgery.* (2017) 3, 110-114.
18. Tantawy AA, Askar SM, Amer HS, Awad A, El-Anwar MW. Hyoid Bone Suspension as a Part of Multilevel Surgery for Obstructive Sleep Apnea Syndrome. *Int Arch Otorhinolaryngol.* 2018 Jul;22(3):266-270. doi: 10.1055/s-0037- 1607227.
19. Yang MC, Hsu YB, Lan MY, Lan MC. The comparison of multilevel surgery (hyoid myotomy and suspension with uvulopalatopharyngoplasty) with CPAP in moderate to severe OSAS patients. *Eur Arch Otorhinolaryngol.* 2020 Aug;277(8):2349-2355. doi: 10.1007/s00405-020-05951-7.
20. Panah ZE, Sharifi A, Zoafa S, Etemadi-Aleagha A, Sohrabpour S, Behzadi M, Teymouri A, Heidari R, Erfanian R. Uvulopalatopharyngoplasty with and without modified thyrohyoid suspension for obstructive sleep apnea treatment: a randomized clinical trial. *Eur Arch Otorhinolaryngol.* 2023 Oct;280(10):4677-4685. doi: 10.1007/s00405-023-08068-9.
21. Van Tassel J, Chio E, Silverman D, Nord RS, Platter D, Abidin MR. Hyoid Suspension With UPPP for the Treatment of Obstructive Sleep Apnea. *Ear Nose Throat J.* 2021 Mar 18:1455613211001132.
22. Handler E, Hamans E, Goldberg AN, Mickelson S. Tongue suspension: An evidence-based review and comparison to hypopharyngeal surgery for OSA. *Laryngoscope.* May 2013; (online) 31. doi:10.1002/lary.24187
23. Sher AE, Schechtman KB, Piccirillo JF. The efficacy of surgical modifications of the upper airway in adults with obstructive sleep apnea syndrome. *Sleep.* 1996 Feb;19(2):156-77. doi: 10.1093/sleep/19.2.156. PMID: 8855039.



International Surgical Sleep Society

1061 East Main Street • Suite 300 East Dundee • Illinois, 60118

[info@surgicalsleeptmeeting.org](mailto:info@surgicalsleeptmeeting.org)